



Data Paper

Dataset on bryophyte species distribution across an elevational gradient on Flores Island

Rosalina Gabriel[‡], Leila N. Morgado[‡], Sílvia Poponessi[§], Débora S. G. Henriques[|],
Márcia C. M. Coelho[‡], Gabriela M. Silveira[¶], Fernando Pereira[‡], Paulo A.V. Borges^{‡#}

[‡] University of Azores, CE3C—Centre for Ecology, Evolution and Environmental Changes, Azorean Biodiversity Group, CHANGE —Global Change and Sustainability Institute, School of Agricultural and Environmental Sciences, Rua Capitão João d'Ávila, Pico da Urze, 9700-042, Angra do Heroísmo, Azores, Portugal

[§] Institute for Alpine Environment, Eurac Research, Viale Druso 1, 39100, Bolzano, Italy

[|] Banco Genético Vegetal Autóctone, Empresa Municipal Cascais Ambiente, Estrada de Vale de Cavalos, 2645-138, Cascais, Portugal

[¶] University of the Azores, Gaspar Frutuoso Foundation, Rua Capitão João D'Ávila, Pico da Urze, São Pedro, 9700-042, Angra do Heroísmo, Portugal

[#] IUCN SSC Monitoring Specialist Group, Angra do Heroísmo, Azores, Portugal

Corresponding author: Paulo A.V. Borges (paulo.av.borges@uac.pt)

Academic editor: Nathaly Guerrero-Ramírez

Received: 19 Jul 2025 | Accepted: 14 Oct 2025 | Published: 19 Nov 2025

Citation: Gabriel R, Morgado LN, Poponessi S, Henriques DSG, Coelho MCM, Silveira GM, Pereira F, Borges PAV (2025) Dataset on bryophyte species distribution across an elevational gradient on Flores Island.

Biodiversity Data Journal 13: e165664. <https://doi.org/10.3897/BDJ.13.e165664>

Abstract

Background

A bryophyte diversity survey was carried out from July 29 to 1 August 2013, in Santa Cruz das Flores, Flores Island (Azores) (39.471185 N Latitude; -31.184692 W Longitude), along an elevational gradient (70, 200, 400, 600 and 800 m a.s.l.). The study employed the Global Island Monitoring Scheme (GIMS) protocol for bryophytes. At each elevation level, three replicates of six substrates colonised by bryophytes (rock, soil, humus, dead wood, tree trunks, leaves) were collected. In total, 385 sampling events generated 1345 species occurrence records, representing 89 bryophyte species (37 mosses; 52 liverworts).

New information

The acrocarpous moss *Fissidens azoricus* (P.de la Varde) Bizot, listed as "Critically Endangered" by the IUCN, had not been reported from Flores Island since 1937; this is the first publication with new locations for the species.

The altitudinal gradient revealed an increase in species richness and abundance with elevation, following a mid-elevation peak pattern, with the highest richness between 400 m a.s.l. (52 species; 364 records) and 600 m a.s.l. (54 species; 402 records), followed by a decline at 800 m a.s.l. (33 species; 148 records). At 70 m a.s.l., 20 species were identified (128 records) and, at 200 m a.s.l., 35 species were recorded (232 records). In terms of substrate preference, the highest species richness and abundance were found colonising epiphytic substrates (58 species; 424 records), followed by terricolous substrates (44 species; 233 records).

Keywords

Azores, bryoflora, elevational gradient, Flores Island, GIMS protocol, liverworts, mosses, MOVECLIM-AZO, natural park, substrates

Introduction

Understanding elevational diversity patterns is essential for guiding conservation actions, predicting the potential community shifts under environmental change scenarios and identifying priority areas for biodiversity protection. Previous studies conducted in the Azores Archipelago have reported high bryophyte species richness on both Pico (Coelho et al. 2021, Gabriel et al. 2024b) and Terceira Islands (Henriques et al. 2017, Coelho et al. 2021, Gabriel et al. 2024a) with richness peaking between 600 and 1000 m a.s.l. This pattern has been largely attributed to the presence of relatively undisturbed native forests at these elevations (Coelho et al. 2016, Henriques et al. 2017, Gabriel et al. 2024a, Gabriel et al. 2024b). In contrast, Flores Island, the westernmost in the Archipelago, had not been intensively surveyed for over two decades. A compilation of literature referencing bryophyte records from Flores Island includes 85 references published between 1844 and 2023, encompassing journal articles, books and book chapters. Notable contributions include Erik Sjögren, Cecilia Sérgio, Pierre and Valentine Allorge and Manuela Sim-Sim (see Suppl. material 1). This data paper aims to address the existing knowledge gap concerning bryophyte diversity on Flores Island.

General description

Purpose: The main objectives of this study were:

1. to conduct a comprehensive survey of bryophytes within the Flores Natural Park, employing the Global Island Monitoring Scheme (GIMS) protocol (Gabriel et al. 2014, Borges et al. 2018), along a transect spanning an elevational gradient from 70 to 800 m a.s.l.;
2. to compile the publications referencing bryophyte records from Flores Island.

Project description

Title: Inventory of bryoflora present in different altitudinal gradients and substrates of Flores Island (Azores)

Personnel: Project conceptualisation: The MOVECLIM project, "Montane vegetation as listening posts for climate change", was originally initiated by the research team led by Claudine Ah-Peng to study and promote tropical bryophytes and ferns as bioindicators of climate change. The project was coordinated in the Azores by Rosalina Gabriel. The monitoring protocol implemented in the project was the GIMS protocol (Ah-Peng et al. 2012, Borges et al. 2018, Gabriel et al. 2024a). This is a standardised method that has also been applied in tropical settings, including La Réunion (Ah-Peng et al. 2012), Madagascar (e.g. Marline et al. (2020)) and other locations.

FIELDWORK:

Site selection and experimental setting: Rosalina Gabriel and Fernando Pereira.

Permits: The Azores Government, through the Environment Department, gave the necessary authorisations to work within the Flores Natural Park. Sample collection: The bryoflora inventory was conducted between 29 July and 1 August 2013, within the Municipality of Santa Cruz das Flores. Sampling was carried out at six sites differing in location, elevation (m a.s.l.) and geographic coordinates (decimal Latitude/Longitude). The fieldwork was coordinated by Rosalina Gabriel, with the participation of Márcia C. M. Coelho, Débora S. G. Henriques and Fernando Pereira.

LAB WORK:

Bryophyte Taxonomic Work: The taxonomic identification of the bryophyte specimens was primarily conducted by Silvia Poponessi, with contributions from Margarida Brito de Azevedo and Rosalina Gabriel. Between 2024 and 2025, a selected subset of specimens was reviewed by Leila N. Morgado and Gabriela M. Silveira, under the supervision of Rosalina Gabriel. Manuela Sim-Sim and Cecília Sérgio helped with the identification of particularly challenging specimens.

MANAGEMENT:

Voucher specimen management: Márcia C. M. Coelho, Débora S. G. Henriques, Rosalina Gabriel.

Database management: Rosalina Gabriel.

Darwin Core databases: Leila N. Morgado, Paulo A. V. Borges, Rosalina Gabriel.

Study area description: Flores Island, one of the nine islands of the Azores Archipelago (Portugal), is situated west of the Mid-Atlantic Ridge, resting on the North American lithospheric plate (Forjaz 2014). Flores is the fourth smallest and the fourth lowest island of the Archipelago, with a land area of 143 km² and a highest elevation of 911 m (Morro Alto). Remnants of well-preserved native vegetation persist, predominantly in the form of *Juniperus brevifolia* forests (Elias et al. 2016) and mire ecosystems (Dias and Melo 2010). While montane forests in the Azores generally occur at elevations between 700 and 1000 m a.s.l., on Flores, they are found at lower altitudes. This shift is attributed to the island's high precipitation levels and the presence of *Sphagnum* spp. (Bryophyta, Sphagnaceae) dominated soils, which provide favourable conditions for the development and maintenance of these forest communities (Elias et al. 2016).

Design description: The bryophyte inventory in the field was conducted from 29 July to 1 August 2013, under the supervision of RG, with field contributions from authors DSGH, MCMC and FP. This fieldwork follows the GIMS protocol for bryophyte sampling (Borges et al. 2018), incorporating both taxonomic information on bryophytes (Ah-Peng et al. 2012) and a suite of environmental variables (Gabriel et al. 2014). Five sites were selected along an altitudinal gradient, spaced at 200-m intervals between 70 m and 800 m above sea level. At each site, two 10 × 10-m plots were established within areas of uniform vegetation structure, with a distance of 10 to 15 m between them. Each plot was subdivided into 25 quadrats (2 m × 2 m), from which three were randomly selected for detailed analysis. From each quadrat, three microplots of 5 cm × 10 cm were collected across six substrate types: rock (rupicolous), soil (terricolous), organic matter (humicolous), decaying wood (lignicolous), bark (epiphytic, at three vertical strata) and leaf/frond surfaces (epiphyllous), whenever possible. Microplots from tree bark were collected in a stratified manner across three vertical zones: low (1–50 cm), medium (51–100 cm) and high (101–200 cm). Specimens were identified in the laboratory to the lowest possible taxonomic level. Additionally, data on cover-abundance and sociability were estimated for each bryophyte taxon within every microplot.

Funding: This study was originally funded by the ERANET BIOME MOVECLIM project "Montane vegetation as listening posts for climate change" (Regional Government of the Azores, grant M2.1.2/F/04/2011/NET). MCMC was supported by the Regional Government of the Azores through FRCT (grant M3.1.2/F/007/2012). RG and PAVB received support from the Azores DRCT Pluriannual Funding (M1.1.A/FUNC.UI&D/010/2021-2024) and FCT (UIDB/00329/2020-2024; DOI: [10.54499/UIDB/00329/2020](https://doi.org/10.54499/UIDB/00329/2020)), under Thematic Line 1 – Integrated ecological assessment of environmental change on

biodiversity. This study is part of the Biodiversa+ project BioMonI – Biodiversity monitoring of island ecosystems, funded by FCT (BiodivMon/0003/2022), which also supported Open Access and provided funding to LNM, RG and PAVB.

Sampling methods

Description: Following the protocol explained above, under 'Design description', each site comprised two 100 m² plots, spaced at 200 m a.s.l., ranging from 70 to 800 m above sea level. The elevational transect extended from Ponta do Ilhéu (70 m a.s.l.) to Morro Alto (800 m a.s.l.). The sampling location names and their corresponding coordinates are listed in Table 1 and illustrated in Fig. 2. At each plot, a general survey of the vascular plant species was performed, noting also the inclination and exposition of the areas and other relevant descriptive details.

Table 1.

Bryophyte sampling sites of the MOVECLIM Project on Flores Island (Azores), surveyed in 2013. The table includes plot codes, locality names, elevations (m a.s.l.), slope, exposure and geographic coordinates (decimal degrees).

| Plot Code | Locality | Elevation (m a.s.l.) | Slope | Exposure | Latitude | Longitude |
|-------------|----------------------------|----------------------|-------|----------|----------|-----------|
| TER_0070_P1 | Ponta do Ilhéu | 70 | 15 | 280 | 39.50633 | -31.19453 |
| TER_0070_P2 | | 77 | 18 | 107 | 39.50622 | -31.19461 |
| TER_0200_P1 | Caminho para Ponta Delgada | 249 | 35 | 141 | 39.50689 | -31.21286 |
| TER_0200_P2 | | 266 | 32 | 14 | 39.50661 | -31.21275 |
| TER_0400_P1 | Outeiros | 399 | 40 | 81 | 39.50192 | -31.20558 |
| TER_0400_P2 | | 399 | 41 | 54 | 39.50183 | -31.20558 |
| TER_0600_P1 | Ribeira do Cascalho | 649 | 5 | 90 | 39.48281 | -31.19042 |
| TER_0600_P2 | | 648 | 8 | 176 | 39.48267 | -31.19033 |
| TER_0800_P1 | Morro Alto | 833 | 14 | 278 | 39.46319 | -31.22594 |
| TER_0800_P2 | | 833 | 15 | 283 | 39.46325 | -31.22600 |

Sampling description: First, each quadrat was carefully observed to check the availability of substrates colonised with bryophytes. Next, microplots of 5 cm x 10 cm, placed on three replicates of the same substrate, were collected with a knife or tweezers to paper bags, identified with the name of the site, altitude, plot, quadrat, substrate and number of the replicate. Other data, such as the phorophyte name and information on evaporation, light, humidity and rugosity of the substrate were also collected.

Quality control: FIELD SAMPLING: Sampling plots were established in environmentally homogeneous zones representing the most characteristic native vegetation at each

surveyed elevation. All collections were conducted by bryologists, who ensured methodological consistency and minimised disturbance by avoiding excessive removal of bryophyte material. **PREPARATION AND STORAGE:** Microplot samples were placed in paper bags and left open in a darkened room until thoroughly dehydrated. Once identification was completed, each sample was transferred to labelled herbarium envelopes. These envelopes are deposited in the Bryophyte Section of the Herbarium of the University of the Azores (AZU), under the designation MOVECLIM – AZORES Project: Bryophytes from Flores Island (2013). **SPECIES IDENTIFICATION (TAXONOMY):** Comprehensive efforts were undertaken to ensure precise taxonomic identification:

1. the most current taxonomic keys and floras were employed under the guidance or direct supervision of experienced bryologists;
2. problematic specimens were submitted to specialists for further assessment and confirmation; and
3. specimens that were extremely small or etiolated were not identified to the species level.




Figure 1. [doi](#)

Map illustrating the grouping of islands in the Azores Archipelago, with Flores Island in the western group.

Moss specimens were identified using the floras of Smith (2004) and Casas et al. (2006), while liverworts were determined based on the works of Paton (1999) and Casas et al. (2009) and the taxonomic key provided by Schumacker and Vána (2005). Identification was further supported by visual references, including Atherton et al. (2010), Lüth (2019) and Elias et al. (2022), alongside digital resources from the British Bryological Society and the Bildatlas der Moose Deutschlands. Nomenclature adhered to Gabriel et al. (2010) and Gabriel et al. (2011) incorporating updates available through the PBA (2025).

DATA REGISTRATION: All identification and metadata were systematically recorded in a structured database in an Excel file. **HERBARIUM DEPOSITION:** Voucher specimens were deposited in the Bryophyte Section of the Herbarium of the University of the Azores (AZU-B), with proper labelling and long-term conservation measures. **DATA STANDARDISATION AND DISSEMINATION:** To ensure data accessibility, these records were formatted following the Darwin Core Archive (DwC-A) standard, facilitating integration into the Global Biodiversity Information Facility (GBIF) and other biodiversity platforms. A core event table was generated, comprising 385 sampling event records, each event corresponding to a single 5 cm × 10 cm microplot. An occurrence extension table was created, including 1345 records that detail all bryophyte taxa identified within the sampled microplots. The standardised data package was published through GBIF and associated biodiversity repositories, contributing to global efforts in bryophyte documentation and conservation research.




Figure 2. [doi](#)

Map of Flores Island showing the bryophyte sampling points surveyed in 2013, in relation to the island's Natural Park boundaries, altimetry and main lakes.

Geographic coverage

Description: The research was carried out on Flores Island, one of the nine islands comprising the Azores Archipelago (Portugal) (Fig. 1). Of the five sampled points, three are located within the limits of the Flores Natural Park, in the Municipality of Santa Cruz das Flores (Fig. 2).

Coordinates: 39.5420 and 39.3470 Latitude; -31.3170 and -31.1020 Longitude.

Taxonomic coverage

Description: The bryophyte specimens collected in this survey comprise representatives of the phyla Bryophyta (mosses) and Marchantiophyta (liverworts). No representatives of the phylum Anthocerotophyta (hornworts) were recorded.

Temporal coverage

Notes: The sampling was performed from 29 July to 1 August 2013.

Collection data

Collection name: MOVECLIM-AZO-FLO_2013_Bryophytes from Flores Island

Collection identifier: 76349556-a70a-4ecc-88a7-cd085b6c875d

Parent collection identifier: AZU_Section Bryophytes

Specimen preservation method: Herbarium preservation

Curatorial unit: Herbarium packet

Usage licence

Usage licence: Creative Commons Public Domain Waiver (CC-Zero)

IP rights notes: Additional data or information related to this study can be obtained by contacting the corresponding author.

Data resources

Data package title: The MOVECLIM – AZORES project: Bryophytes from Flores Island (2013).

Resource link: http://ipt.gbif.pt/ipt/resource?r=moveclim_azores_flores_island_2012

Alternative identifiers: <https://www.gbif.org/dataset/f846b120-a1a7-4b1d-a4b5-e0baf6f46775>

Number of data sets: 2

Data set name: Event Table

Character set: UTF-8

Download URL: http://ipt.gbif.pt/ipt/resource?r=moveclim_azores_flores_island_2012

Data format: Darwin Core Archive**Data format version: V1.1**

Description: The dataset was published in the Global Biodiversity Information Facility platform, GBIF (Gabriel et al. 2025). The following data table includes all the records for which a taxonomic identification of the species was possible. The dataset submitted to GBIF is structured as a sample event dataset that has been published as a Darwin Core Archive (DwCA), which is a standardised format for sharing biodiversity data as a set of one or more data tables. The core data file contains 385 records (eventID). This GBIF IPT (Integrated Publishing Toolkit, Version 2.5.6) archives the data and, thus, serves as the data repository. The data and resource metadata are available for download in the Portuguese GBIF Portal IPT (Gabriel et al. 2025).

| Column label | Column description |
|---------------------|--|
| eventID | Identifier of the events, unique for the dataset. |
| type | Type of the record, as defined by the Dublin Core Standard. |
| datasetName | Name of the dataset that in current projects is "MOVECLIM-AZO_2013_Bryophytes from Flores Island". |
| samplingProtocol | The sampling protocol used to collect the species. |
| eventDate | Range during which the record was collected. |
| day | The day of the month on which the Event occurred. |
| month | The month in which the Event occurred. |
| year | The year in which the Event occurred. |
| habitat | The habitat for an Event. |
| continent | The name of the continent in which the Location occurs (Europe). |
| islandGroup | Name of archipelago, always Azores in the dataset. |
| island | Name of the island, always Flores in the dataset. |
| country | Country of the sampling site, always Portugal in the dataset. |
| countryCode | ISO code of the country of the sampling site, always PT in the dataset. |
| municipality | Municipality of the sampling site, always Santa Cruz das Flores in the dataset. |
| locality | The specific name of the locality. |
| verbatimElevation | The original description of the elevation (altitude above sea level in metres) of the location. |
| verbatimCoordinates | Original coordinates recorded. |
| decimalLatitude | Approximate decimal latitude. |

| | |
|-------------------------------|---|
| decimalLongitude | Approximate decimal longitude. |
| geodeticDatum | Standard Global Positioning System coordinate reference for the location of the sample collection points. |
| coordinateUncertaintyInMetres | Uncertain value of coordinate metrics. |
| coordinatePrecision | Value in decimal degrees to a precision of five decimal places. |
| georeferenceSources | Navigation system used to record the location of sample collections. |

Data set name: Occurrence table

Character set: UTF-8

Download URL: http://ipt.gbif.pt/ipt/resource?r=moveclim_azores_flores_island_2012

Data format: Darwin Core Archive format

Data format version: V1.1

Description: The dataset was published in the Global Biodiversity Information Facility platform, GBIF (Gabriel et al. 2025). The following data table includes all the records for which a taxonomic identification of the species was possible. The dataset submitted to GBIF is structured as an occurrence table that has been published as a Darwin Core Archive (DwCA), which is a standardised format for sharing biodiversity data as a set of one or more data tables. The core data file contains 1345 records (occurrenceID). This GBIF IPT (Integrated Publishing Toolkit, Version 2.5.6) archives the data and, thus, serves as the data repository. The data and resource metadata are available for download in the Portuguese GBIF Portal IPT (Gabriel et al. 2025).

| Column label | Column description |
|-------------------|--|
| eventID | Identifier of the events, unique for the dataset. |
| licence | Reference to the licence under which the record is published. |
| institutionID | The identity of the institution publishing the data. |
| institutionCode | The code of the institution publishing the data. |
| collectionID | Identifier of the collection, unique for each specimens are conserved. |
| collectionCode | The code of the collection where the specimens are conserved. |
| datasetName | Project reference. |
| type | Characteristics of the object of study. |
| basisOfRecord | The nature of the data record. |
| dynamicProperties | A list of additional measurements, facts, characteristics or assertions about the record, including IUCN categories and colonisation status of taxa. |
| occurrenceID | Identifier of the record, coded as a global unique identifier. |

| | |
|--------------------------|---|
| recordNumber | An identifier given to the Occurrence at the time it was recorded. |
| recordedBy | A list (concatenated and separated) of names of people, groups or organisations responsible for recording the original Occurrence. |
| identifiedBy | A list (concatenated and separated) of names of people, who made the identification. |
| dateIdentified | Date of species identification. |
| disposition | The current state of a specimen with respect to the collection identified in collectionCode or collectionID. |
| taxonRank | Lowest taxonomic rank of the record. |
| kingdom | Kingdom name. |
| phylum | Phylum name. |
| class | Class name. |
| order | Order name. |
| family | Family name. |
| genus | Genus name. |
| specificEpithet | Specific epithet. |
| infraspecificEpithet | Infraspecific epithet at subspecies level. |
| scientificNameAuthorship | The authorship information for the scientificName formatted according to the conventions of the applicable nomenclaturalCode. |
| ScientificName | Complete scientific name including author. |
| organismQuantity | A number or enumeration value for the quantity of organisms (i, solitary specimen - one or few individuals; p, occasional and less than 5% cover; 1, less than 5% cover of total area; 2, 5%-25% of total area; 3, 25%-50% of total area; 4, 50%-75% of total area; 5, 75%-100% of total area). |
| organismQuantityType | Braun-Blanquet Scale. |
| establishmentMeans | The process of establishment of the species in the location, using a controlled vocabulary: 'Azores endemic', 'European endemic', 'Macaronesian endemic', 'native'. |
| occurrenceRemarks | Remarks on the material or surface where the biological specimen was collected. |

Additional information

This inventory of the bryoflora of Flores Island (Azores) recorded 385 sampling events, yielding 1345 identified specimens, representing 37 families, 58 genera and 89 species, with a average number of species per sample of 3.26.

Of all the specimens collected, only 5.4% could not be identified to species level, a relatively low proportion, indicating a high level of taxonomic resolution.

The phylum Marchantiophyta (liverworts) was the most represented, accounting for 52 species (58.43%), followed by phylum Bryophyta (mosses) with 37 species (41.57%). No representatives of the phylum Anthocerotophyta (hornworts) were collected during this study.

In terms of colonisation status, most species were classified as native (n = 66). Additionally, 23 endemic species were identified, with varying geographic ranges: 12 are considered European endemics, seven are Macaronesian endemics and four are endemic to the Azores. No invasive bryophytes were found within the sampled sites.

According to the IUCN Red List (Hodgetts et al. 2019), one species is considered Critically Endangered (*Fissidens azoricus*), four species are considered Endangered (*Echinodium renauldii*, *Calypogeia azorica*, *Cololejeunea sintenisii*, *Leptoscyphus porphyrius* subsp. *azoricus*), another four species are considered Vulnerable (*Isothecium prolixum*, *Andoa berthelotiana*, *Cololejeunea azorica*, *Lejeunea mandonii*), 14 species are considered Near Threatened, while 64 species are classified as Least Concern (Table 2). Two species are not evaluated: *Daltonia lindigiana*, an American moss species restricted to the Azores in Europe and *Hypnum resupinatum*, sometimes considered part of the *Hypnum cupressiforme* complex (Hodgetts et al. 2020).

| Table 2. List of species and subspecies sampled on Flores Island, with their colonisation status categories (Azorean endemic, European endemic, Macaronesian endemic and Native) and IUCN status (Critically Endangered, Endangered, Least Concern, Near Threatened, Vulnerable and Not Evaluated). | | | |
|--|--|----------------------|-----------------------|
| Phylum | Scientific Name | Colonisation status | IUCN Status |
| Bryophyta | <i>Andoa berthelotiana</i> (Mont.) Ochyra | Macaronesian endemic | Vulnerable |
| | <i>Atrichum undulatum</i> (Hedw.) P.Beauv. | Native | Least concern |
| | <i>Brachythecium mildeanum</i> (Schimp.) Schimp. | Native | Least concern |
| | <i>Campylopus flexuosus</i> (Hedw.) Brid. | Native | Least concern |
| | <i>Campylopus shawii</i> Wilson | Native | Least concern |
| | <i>Daltonia lindigiana</i> Hampe | Native | Not evaluated |
| | <i>Echinodium renauldii</i> (Cardot) Broth. | Azorean endemic | Endangered |
| | <i>Exsertotheca crispa</i> (Hedw.) S.Olsson, Enroth & D.Quandt | Native | Least concern |
| | <i>Fissidens adianthoides</i> Hedw. | Native | Least concern |
| | <i>Fissidens azoricus</i> (P.de la Varde) Bizot | Azorean endemic | Critically endangered |
| | <i>Fissidens crispus</i> Mont. | Native | Least concern |

| Phylum | Scientific Name | Colonisation status | IUCN Status |
|-----------------|--|----------------------|-----------------|
| | <i>Fissidens polyphyllus</i> Wilson ex Bruch & Schimp. | Native | Least concern |
| | <i>Fissidens serrulatus</i> Brid. | Native | Least concern |
| | <i>Fissidens taxifolius</i> Hedw. | Native | Least concern |
| | <i>Heterocladium flaccidum</i> (Schimp.) A.J.E.Sm. | Native | Least concern |
| | <i>Heterocladium wulfsbergii</i> I.Hagen | Native | Least concern |
| | <i>Hypnum jutlandicum</i> Holmen & E.Warncke | Native | Least concern |
| | <i>Hypnum resupinatum</i> Taylor | Native | Not evaluated |
| | <i>Hypnum uncinulatum</i> Jur. | European endemic | Least concern |
| | <i>Isopterygiopsis pulchella</i> (Hedw.) Z.Iwats. | Native | Near threatened |
| | <i>Isothecium prolixum</i> (Mitt.) M.Stech, Sim-Sim, Tangney & D.Quandt | Macaronesian endemic | Vulnerable |
| | <i>Kindbergia praelonga</i> (Hedw.) Ochyra | Native | Least concern |
| | <i>Leucobryum glaucum</i> (Hedw.) Ångstr. | Native | Least concern |
| | <i>Leucobryum juniperoideum</i> (Brid.) Müll.Hal. | Native | Least concern |
| | <i>Mnium hornum</i> Hedw. | Native | Least concern |
| | <i>Nogopterium gracile</i> (Hedw.) Crosby & W.R.Buck | Native | Least concern |
| | <i>Polytrichum commune</i> Hedw. | Native | Least concern |
| | <i>Pseudoamblystegium subtile</i> (Hedw.) Vanderp. & Hedenäs | Native | Least concern |
| | <i>Pseudotaxiphyllum laetevirens</i> (Dixon & Luisier ex F.Koppe & Düll) Hedenäs | European endemic | Near threatened |
| | <i>Rhynchostegiella azorica</i> Hedenäs & Vanderp. | Azorean endemic | Near threatened |
| | <i>Sematophyllum substrumulosum</i> (Hampe) E.Britton | Native | Least concern |
| | <i>Sphagnum palustre</i> L. | Native | Least concern |
| | <i>Sphagnum papillosum</i> Lindb. | Native | Least concern |
| | <i>Tetastichium virens</i> (Cardot) S.P.Churchill | Macaronesian endemic | Near threatened |
| | <i>Thamnobryum maderense</i> (Kindb.) Hedenäs | Native | Near threatened |
| | <i>Thuidium tamariscinum</i> (Hedw.) Schimp. | Native | Least concern |
| | <i>Trichostomum brachydontium</i> Bruch | Native | Least concern |
| Marchantiophyta | <i>Calypogeia arguta</i> Nees & Mont. | Native | Least concern |
| | <i>Calypogeia azorica</i> Bischl. | Macaronesian endemic | Endangered |

| Phylum | Scientific Name | Colonisation status | IUCN Status |
|--------|---|----------------------|-----------------|
| | <i>Calypogeia fissa</i> (L.) Raddi | Native | Least concern |
| | <i>Calypogeia muelleriana</i> (Schiffn.) Müll.Frib. | Native | Least concern |
| | <i>Cephalozia bicuspidata</i> (L.) Dumort. | Native | Least concern |
| | <i>Chiloscyphus polyanthos</i> (L.) Corda | Native | Least concern |
| | <i>Cololejeunea azorica</i> V.Allorge & Jovet-Ast | Native | Vulnerable |
| | <i>Cololejeunea microscopica</i> (Taylor) Schiffn. | Native | Least concern |
| | <i>Cololejeunea sintenisii</i> (Steph.) Pócs | Native | Endangered |
| | <i>Colura calyptrifolia</i> (Hook.) Dumort. | Native | Least concern |
| | <i>Drepanolejeunea hamatifolia</i> (Hook.) Schiffn. | Native | Least concern |
| | <i>Frullania acicularis</i> Hentschel & von Konrat | Native | Near threatened |
| | <i>Frullania azorica</i> Sim-Sim, Sérgio, Mues & Kraut | European endemic | Least concern |
| | <i>Frullania microphylla</i> (Gottsche) Pearson | European endemic | Least concern |
| | <i>Frullania teneriffae</i> (F.Weber) Nees | European endemic | Least concern |
| | <i>Fuscocephaloziopsis crassifolia</i> (Lindenb. & Gottsche) Váňa & L.Söderstr. | Native | Least concern |
| | <i>Fuscocephaloziopsis lunulifolia</i> (Dumort.) Váňa & L.Söderstr. | Native | Least concern |
| | <i>Geocalyx graveolens</i> (Schrad.) Nees | Native | Near threatened |
| | <i>Harpalejeunea mollerii</i> (Steph.) Grolle | Native | Least concern |
| | <i>Heteroscyphus denticulatus</i> (Mitt.) Schiffn. | Macaronesian endemic | Near threatened |
| | <i>Isopaches bicrenatus</i> (Schmidel ex Hoffm.) H.Buch | Native | Least concern |
| | <i>Jubula hutchinsiae</i> (Hook.) Dumort. | Native | Least concern |
| | <i>Lejeunea cavifolia</i> (Ehrh.) Lindb. | Native | Least concern |
| | <i>Lejeunea eckloniana</i> Lindenb. | Native | Least concern |
| | <i>Lejeunea flava</i> (Sw.) Nees | Native | Near threatened |
| | <i>Lejeunea hibernica</i> Bischl., H.A.Mill. & Bonner ex Grolle | European endemic | Near threatened |
| | <i>Lejeunea lamacerina</i> (Steph.) Schiffn. | Native | Least concern |
| | <i>Lejeunea mandonii</i> (Steph.) Müll.Frib. | Native | Vulnerable |
| | <i>Lejeunea patens</i> Lindb. | Native | Least concern |
| | <i>Lepidozia reptans</i> (L.) Dumort. | Native | Least concern |
| | <i>Leptoscyphus cuneifolius</i> (Hook.) Mitt. | Native | Least concern |

| Phylum | Scientific Name | Colonisation status | IUCN Status |
|--------|--|----------------------|-----------------|
| | <i>Leptoscyphus porphyrius</i> subsp. <i>azoricus</i> (H.Buch & Perss.) Vanderp. & Heinrichs | Azorean endemic | Endangered |
| | <i>Lophocolea bidentata</i> (L.) Dumort. | Native | Least concern |
| | <i>Lophocolea fragrans</i> (Moris & De Not.) Gottsche, Lindenb. & Nees | Native | Least concern |
| | <i>Marchesinia mackaii</i> (Hook.) Gray | European endemic | Least concern |
| | <i>Metzgeria furcata</i> (L.) Corda | Native | Least concern |
| | <i>Microlejeunea ulicina</i> (Taylor) Steph. | Native | Least concern |
| | <i>Myriocoleopsis minutissima</i> (Sm.) R.L. Zhu, Y.Yu & Pócs | Native | Least concern |
| | <i>Nowellia curvifolia</i> (Dicks.) Mitt. | Native | Least concern |
| | <i>Plagiochila bifaria</i> (Sw.) Lindenb. | Native | Least concern |
| | <i>Porella canariensis</i> (F.Weber) Underw. | European endemic | Least concern |
| | <i>Porella obtusata</i> (Taylor) Trevis. | Native | Least concern |
| | <i>Radula aquilegia</i> (Hook.f. & Taylor) Gottsche, Lindenb. & Nees | European endemic | Least concern |
| | <i>Radula carringtonii</i> J.B. Jack | European endemic | Near threatened |
| | <i>Radula holzii</i> Spruce | European endemic | Near threatened |
| | <i>Radula wichurae</i> Steph. | Macaronesian endemic | Near threatened |
| | <i>Riccardia chamedryfolia</i> (With.) Grolle | Native | Least concern |
| | <i>Saccogyna viticulosa</i> (L.) Dumort. | European endemic | Least concern |
| | <i>Scapania gracilis</i> Lindb. | Native | Least concern |
| | <i>Scapania nemorea</i> (L.) Grolle | Native | Least concern |
| | <i>Telaranea azorica</i> (H.Buch & Perss.) Pócs | Macaronesian endemic | Near threatened |
| | <i>Telaranea europaea</i> J.J. Engel & G.L. Merr. | Native | Least concern |

Additionally, two Azorean endemics and one Macaronesian endemic are classified as Endangered and two are listed as Vulnerable, according to the IUCN Red List (Table 2). Notably, this study recorded the presence of *Fissidens azoricus*, a species listed as Critically Endangered by the IUCN and not recorded on Flores Island since 1937. This emphasises the importance of bryophyte inventory efforts for conservation and biodiversity monitoring.

Acknowledgements

The authors are deeply grateful to Dr. Cecília Sérgio and Prof. Manuela Sim-Sim for their help with the identification of challenging samples of bryophytes and to all the funding entities that allowed the execution of the present study. This study is part of the Biodiversa+ project BioMonI – Biodiversity monitoring of island ecosystems, funded by FCT (BiodivMon/0003/2022), which also supported Open Access and provided funding to LNM, RG and PAVB.

Author contributions

Rosalina Gabriel: Conceptualisation; Methodology; Research (laboratory work); Resources; Data Curation; Darwin Core dataset preparation; Formal analysis and interpretation; Manuscript writing.

Leila Nunes Morgado: Formal analysis and interpretation; Research (laboratory work); Manuscript writing.

Débora S. G. Henriques, Márcia C. M. Coelho: Research (field and laboratory work).

Fernando Pereira: Research (field work).

Silvia Poponessi, Gabriela Melo Silveira: Research (laboratory work).

Paulo A. V. Borges: Darwin Core dataset revision, GBIF IPT management.

References

- Ah-Peng C, Wilding N, Kluge J, Descamps-Julien B, Bardat J, Chuah-Petiot M, Strasberg D, Hedderson TJ (2012) Bryophyte diversity and range size distribution along two altitudinal gradients: Continent vs. island. *Acta Oecologica* 42: 58-65. <https://doi.org/10.1016/j.actao.2012.04.010>
- Atherton I, Bosanquet S, Lawley M (Eds) (2010) Mosses and liverworts of Britain and Ireland: a field guide. Plymouth: British Bryological Society [ISBN 978-0-9561310-1-0]
- Borges PAV, Cardoso P, Kreft H, Whittaker R, Fattorini S, Emerson B, Gil A, Gillespie R, Matthews T, Santos AC, Steinbauer M, Thébaud C, Ah-Peng C, Amorim I, Aranda SC, Arroz AM, Azevedo J, Boeiro M, Borda-de-Água L, Carvalho JC, Elias R, Fernández-Palacios JM, Florencio M, González-Mancebo J, Heaney L, Hortal J, Kueffer C, Lequette B, Martín-Esquivel JL, López H, Lamelas-López L, Marcelino J, Nunes R, Oromí P, Patiño J, Pérez A, Rego C, Ribeiro S, Rigal F, Rodrigues P, Rominger A, Santos-Reis M, Schaefer H, Sérgio C, Serrano AM, Sim-Sim M, Stephenson PJ, Soares A, Strasberg D, Vanderporten A, Vieira V, Gabriel R (2018) Global Island Monitoring Scheme (GIMS): a proposal for the long-term coordinated survey and monitoring of native island forest biota. *Biodiversity and Conservation* 27 (10): 2567-2586. <https://doi.org/10.1007/s10531-018-1553-7>

- Casas C, Brugués M, Cros RM, Sérgio C (2006) Handbook of mosses of the Iberian Peninsula and the Balearic Islands: illustrated keys to genera and species. Institut d'Estudis Catalans [ISBN 84-7283-865-X]
- Casas C, Brugués M, Cros RM, Sérgio C, Infante M (2009) Handbook of liverworts and hornworts of the Iberian Peninsula and the Balearic Islands: illustrated keys to genera and species. Institut d'estudis Catalans [ISBN 9788492583553]
- Coelho MC, Elias RB, Kluge J, Pereira F, Henriques DSG, Aranda SC, Borges PAV, Ah-Peng C, Gabriel R (2016) Vascular plants on Pico island transect. Long-term monitoring across elevational gradients (II): vascular plants on Pico Island (Azores) transect. *Arquipelago - Life and Marine Sciences* 33: 21-44.
- Coelho MM, Gabriel R, Hespanhol H, Borges PV, Ah-Peng C (2021) Bryophyte Diversity along an Elevational Gradient on Pico Island (Azores, Portugal). *Diversity* 13 (4). <https://doi.org/10.3390/d13040162>
- Dias E, Melo C (2010) Factors influencing the distribution of Azorean mountain vegetation: implications for nature conservation. *Biodiversity and Conservation* 19 (12): 3311-3326. <https://doi.org/10.1007/s10531-010-9894-x>
- Elias R, Gil A, Silva L, Fernández-Palacios J, Azevedo E, Reis F (2016) Natural zonal vegetation of the Azores Islands: characterization and potential distribution. *Phytocoenologia* 46 (2): 107-123. <https://doi.org/10.1127/phyto/2016/0132>
- Elias RB, Rodrigues AFF, Gabriel R (2022) Guia prático da flora nativa dos Açores/Field Guide of Azorean Native Flora. 1, 1. Instituto Açoriano de Cultura, Angra do Heroísmo. [ISBN 9789898225740]
- Forjaz VH (2014) Atlas básico dos Açores. OVGA, Ponta Delgada.
- Gabriel R, Sjögren E, Schumacker R, Sérgio C, Aranda S, Claro D, Homem N, Martins B (2010) List of bryophytes (Anthocerotophyta, Marchantiophyta, Bryophyta). In: Borges PAV, Costa A, Cunha R, Gabriel R, et al. (Eds) A List of the Terrestrial and Marine Biota from the Azores. 1. Princípiia, Cascais. [ISBN 978-989-8131-75-1].
- Gabriel R, Homem N, Couto A, Aranda SC, Borges PAV (2011) Azorean Bryophytes: A preliminary review of rarity patterns. *Açoreana* 7: 149-206.
- Gabriel R, Coelho MCM, Henriques DSG, Borges PAV, Elias RB, Kluge J, Ah-Peng C (2014) Long-term monitoring across elevational gradients to assess ecological hypothesis: a description of standardized sampling methods in oceanic islands and first results Arquipelago. *Arquipelago - Life and Marine Sciences* 31: 45-57. URL: https://repositorio.uac.pt/bitstream/10400.3/3296/1/LMSpp45-67_Gabriel_et_al_N31.pdf
- Gabriel R, Morgado L, Henriques D, Coelho M, Hernández-Hernández R, Borges P (2024a) The MOVECLIM – AZORES project: Bryophytes from Terceira Island along an elevation gradient. *Biodiversity Data Journal* 12: e131935. <https://doi.org/10.3897/bdj.12.e131935>
- Gabriel R, Morgado L, Borges P, Coelho M, Aranda S, Henriques D, Sérgio C, Hespanhol H, Pereira F, Sim-Sim M, Ah-Peng C (2024b) The MOVECLIM – AZORES project: Bryophytes from Pico Island along an elevation gradient. *Biodiversity Data Journal* 12 "30" href="https://doi.org/10.3897/bdj.12.e117890" target=" blank"> <https://doi.org/10.3897/bdj.12.e117890>
- Gabriel R, Morgado LN, Poponessi S, Henriques DSG, Coelho MCM, Silveira GM, Pereira F, Borges PAV (2025) The MOVECLIM – AZORES project: Bryophytes from Flores Island (2013). 1.1. GBIF. Release date: 2025-7-18. URL: <https://www.gbif.org/dataset/f846b120-a1a7-4b1d-a4b5-e0baf6f46775>

- Henriques DS, Elias RB, Coelho MC, Hernández RH, Pereira F, Gabriel R (2017) Long-term monitoring across elevational gradients (III): Vascular plants on Terceira Island (Azores) transect. *Arquipelago - Life and Marine Sciences* 34: 1-20.
- Henriques DSG, Borges PAV, Gabriel R (2017) Regional processes drive bryophyte diversity and community composition in a small oceanic island. *Community Ecology* 18 (2): 193-202. <https://doi.org/10.1556/168.2017.18.2.9>
- Hodgetts N, Calix M, Englefield E, Fettes N, García Criado M, Patin L, et al. (2019) A miniature world in decline: European Red List of mosses, liverworts and hornworts. IUCN, Brussels, Belgium, 620 pp. [ISBN **0873-4704**] "30" [href="https://doi.org/10.2305/IUCN.CH.2019.ERL.2.en" target=" blank"> https://doi.org/10.2305/IUCN.CH.2019.ERL.2.en](https://doi.org/10.2305/IUCN.CH.2019.ERL.2.en)
- Hodgetts NG, Söderström L, Blockeel TL, Caspari S, Ignatov MS, Konstantinova NA, Lockhart N, Papp B, Schröck C, Sim-Sim M, Bell D, Bell NE, Blom HH, Bruggeman-Nannenga MA, Brugués M, Enroth J, Flatberg KI, Garilleti R, Hedenäs L, Holyoak DT, Hugonnot V, Kariyawasam I, Köckinger H, Kučera J, Lara F, Porley RD (2020) An annotated checklist of bryophytes of Europe, Macaronesia and Cyprus. *Journal of Bryology* 42 (1): 1-116. <https://doi.org/10.1080/03736687.2019.1694329>
- Lüth M (2019) Mosses of Europe - A photographic flora. Freiburg [ISBN 978-3-00-062952-5]
- Marline L, Ah-Peng C, Hedderson TJ (2020) Epiphytic bryophyte diversity and range distributions along an elevational gradient in Marojejy, Madagascar. *Biotropica* 52 (4): 616-626. <https://doi.org/10.1111/btp.12781>
- Paton JA (1999) The liverwort flora of the British Isles. Harley Books [ISBN **0946589607**] "30" [href="https://doi.org/10.1163/9789004285385" target=" blank"> https://doi.org/10.1163/9789004285385](https://doi.org/10.1163/9789004285385)
- PBA (2025) Portal da Biodiversidade dos Açores | Azorean Biodiversity Group. <https://azoresbiportal.uac.pt>. Accessed on: 2025-6-03.
- Schumacker R, Vána J (2005) Identification keys to the liverworts and hornworts of Europe and Macaronesia. 2nd Edition. Station Scientifique des Hautes-Fagnes
- Smith AJE (2004) The moss flora of Britain and Ireland. Cambridge university press [ISBN **9780511541858**] "30" [href="https://doi.org/10.1017/CBO9780511541858" target=" blank"> https://doi.org/10.1017/CBO9780511541858](https://doi.org/10.1017/CBO9780511541858)

Supplementary material

Suppl. material 1: Compilation of Publications [doi](#)

Authors: Rosalina Gabriel

Data type: Compilation of Publications

Brief description: List of Publications on Bryophytes in Flores Island (Azores) from 1844 to 2023.

[Download file](#) (21.42 kb)